



VÕITLUS BIOKILMIDEGA

MIKS TEIE ANTIBIOOTIKUD ANO SEENEVASTASED F AIL

Lahendused Lyme'i tõve, kroonilise sinusiidi,
Kopsupõletik, pärmseente infektsioonid, haavad, kõrvad
Infektsioonid, igemehaigused, soolehaigused,
Halb hingeõhk, tsüstiline fibroos ja implantaadid


KROONILISE HAIGUSE MÕISTLUSES PÕHINE PUUDUV OSAK

JAMES SCHALLER, M.D., M.A.R.

KIMBERLY MOUNTJOY, M.S.

Mis on biofilm?

Lihne, teaduslik biokile määratlus: mis tahes mikroorganismide rühm, mille rakud kleepuvad pinnal üksteise külge. Need asuvad tavaliselt nende loodud kihis, mida nimetatakse limadeks.



Võrrelge biokilet praemunaga. Praemuna keskel olev kollane munakollane on bakteriaalne või seeninfektsioon.

Suuremat valget osa, mis munakollast ümbritseb, võib nimetada "biokileks". See kaitseb sisemist infektsiooni ehk munakollast nii antibiootikumide kui ka inimese immuunsüsteemi eest.

Muna välisserval on mõned väga väikesed praetud servad. Neid on muna suuruse tõttu lihtne vahele jätta. Me teeskleme, et need on antibiootikumid või infektsioone hävitavad kemikaalid. Need on kasutud, kuna ei pääse kunagi muna valgest välisservast mööda. Munavalge on neile nagu sein.

Kellel on biofilmiinfektsioonid?

Kui õpite tundma tohutut mitmekesisust asukohtade ja olukordade kohta, kus biofilmid on levinud, ja arvate, et see on sageli bakterite ja seenorganismide tavapärane seisund, hakkate mõistma, et kellelgi võib olla biokileinfektsioon või -infektsioonid.

Mida me sellest raamatust otsime?

Järgmises materjalis näidatakse mitmeid viise, kuidas "munavalgest" ehk biokilest läbi murda. Kui see juhtub, on tavaliselt palju lihtsam munakollase või kollase keskuse poolt kujutatud nakkus

Biofilmid on kannatuste ja surma peamine põhjus

Biofilmi keha asukohad ja olukorrad

- üle 2 nädala kestev infektsioon
- Alla 6-aastaste laste peamine surmapõhjus
- Hambakatt – inimese suus on umbes 25 000 liiki baktereid, millest umbes 1000 asub hambakatu biokiles.

- Pärmseente infektsioonid
- Postkirurgilised infektsioonid
- Vähk
- Halb hingeõhk
- Igemehaigus või periodontiit*
- Hammaste lagunemine • Kopsuinfektsioonid
- Kuseteede infektsioonid
- Suubakterid – võivad kahjustada südameartereid ja põhjustada surma ning suurendada soolevähi teket
- Kroonilised kõrvapõletikud
- Siinusinfektsioonid**
- Krooniline tonsilliit
- Haavad

- Hambaharjapead – sealhulgas heliliselt liikuvad pead

- kateetrid uriini eemaldamiseks
- Kunstlikud põlved, puusad ja muud asendused
- Südameklapi infektsioonid
- Kahjustused või haavandid
- Lyme'i tõbi
- Igat tüüpi IV kateetrid
- Kuseteede kateetrid
- Kontaktläätсед
- Implanteeritud seadmed – iga implanteeritud või sisestatud seade võib saata baktereid ajju, maksa või neerudesse.
- Kroonilised eesnäärmeinfektsioonid
- Leegionäride haigus ja paljud teised biotoksiinibakterid, mis plahvatavad igas siseveekogus
- Hallitushaigused – mis võivad tekkida hallituse kogunemisest mistahes seisvas siseveekogus, nt üleujutus, katuse-, keldri- või aknalekked, õhuniisutajad, kasutamata Waterpik™ või muud hambapuhastusseadmed,

kondenseerumine vahelduvvoolu kanalitesse jne. • Tsüstiline fibroos – liigne lima tootmine hingamisteedes võimaldab bakteritel nagu *Pseudomonas aeruginosa* võita biokilekatte taga bakterite tapjaid.

- Kaotatud kehaosad
- Naha, juuste või küünte infektsioonid
- Artriit
- Endokardiit
- Luuinfektsioonid
- Vinnid

Loendisse võiks lisada palju muid asju, sealhulgas sügavalt tõsiseid probleeme biokile saastumisega vees ja kümneid muid tervisega seotud ja tootmistavasid.

*Pensionil olev hambaarst doktor David Kennedy kahetses, et enamikul täiskasvanud ameeriklastel on igemehaigus – veel üks bakteriaalne biokile haigus, millega kaasneb krooniline infektsioon. Niisiis, kui laialt levinud on see vargsi tervishoiuepideemia?

**Ondine Biopharma intervjuust [Richard Longlandiga] selgus, et 38 000 000 inimesel selles riigis on (või oli) krooniline põsekoopaprobleem.

***Ricardo Murga; Terri S. Forster. Biokilede roll *Legionella pneumophila* ellujäämisel joogivee mudelis. *Microbiology* (2001), 147, 3121–3126.

VÕITLUS BOKILMIDEGA

Miks teie antibiootikumid ja seenevastased ravimid ebaõnnestuvad

**Lahendused Lyme'i tõve, kroonilise sinusiidi,
Kopsupõletik, pärmseente infektsioonid, haavad, kõrvad
Infektsioonid, igemehaigused, soolehaigused,
Halb hingeõhk, tsüstiline fibroos ja implantaadid**

Kroonilise haiguse pusle peamine kadunud tük

James Schaller, MD, MAR

ja

Kimberly Mountjoy, MS

**Rahvusvaheline nakkushaiguste ajakirjandus
Bank Towers • Newgate Center (sviit 305)
5150 Tamiami Trail North [Highway 41]
Napoli, Florida 34103**

Praeguste biofilmide elupäästvate vastuste tegemine Selge ja kivikindel

Praegu võite lugeda kahe aasta väärtuses biofilmi hävitamise võimalusi lehtedest, ajaveebidest ja raamatutest. Selleks kulub teil 1000–1500 tundi. Ja teil oleks välja pakkuda mitu võimalust. Siin on mõned näited valikutest, mida leiate nendest paberitest, ajaveebidest ja raamatutest.

Vältige magneesiumi	EDTA	Mesilaspiim
Vältige suhkruid ja teravilju	DMSO	tüümian
NAC	Vankomütsiin	Sidrunhein
Norspermiidiin	Gentamütsiin	Serrapeptidaas
Cis 2 - detseenhape	Banderol	2-aminobensimidiasool
Lumbrokinaas	Vältige rasvu	Ehhinokandiinid

Kuidas leida mõistlikku turundust ja usaldust biofilmi agendi kui lahenduse vastu?

Tom ja Lisa blogivad, et toode "x" ja retsept "d" on erakordsed ravimeetodid kroonilise väsimuse (CFS) ja fibromüalgia (FM) biokileinfektsioonide kahjustamiseks. Inimesed on põnevil, kuna nende tavaarstil pole suurt lahendust ega huvi biokileinfektsioonide vastu.

Probleem on selles, et "x" või "d" võivad olla kasulikud biokile kahjustamiseks või haigusest ülesaamiseks. Kuid olge kiirete linkide loomisel ettevaatlik. Ravi "a" võib toimida ainult kümne infektsiooni biofilmis ja meil on tõendeid selle kohta, et see toimib ainult kolme infektsiooni korral.

Meie eesmärk on näidata teile, mida näitavad head uuringud, et teie ja teie arst saaksite alustada faktidest ja mõistaksite võimalike biofilmikatsetuste põhjuseid.

Näiteks võib teie infektsioon raua kasutamise poolest sarnaneda Lyme'iga. Saito ja paljud teised teatavad, et erinevalt kõigist teistest teadaolevatest organismidest võib borrelia, Lyme'i tõve põhjustaja, eksisteerida ilma rauata, metallita, mida kogu muu elu vajab. Selle asemel kasutab Borrelia mangaani.

Mis siis, kui leitakse, et teie biokilepõhisel haigusel on tulevikus sama võime elada hästi ilma rauata? See võib tähendada, et Lyme'i tõve biokile õõnestav biokile võib teie jaoks sobida. Bakterite ja seeninfektsioonide biokilel on biokile lõhkuja suhtes sarnane haavatavus. Teie nakkuse toimimise teadmine võib aidata kindlaks teha, milline biokile aine töötab.

<http://phys.org/news/2013-03-scientists-reveal-quirky-feature-lyme.html#jCp>.
Vaadatud 26. märtsil 2014.

Sisu

Meditsiini revolutsioon	1
Biofilmide viis etappi	2
Sissejuhatavad biofilmi pildid	3
Üldine bakter, mida leidub kogu inimese nahal	4
Staph-nakkused on muutunud enamiku antibiootikumide suhtes resistentseks	5
Meditsiiniringkond ignoreerib biofilme	20
"Biofilmide" selgeks tegemine	21
Kannatuste, madalama funktsionaalsuse, puude ja surma ennetamine biofilmide tõttu	24
Biokile eemaldamine muudab antibiootikumid tõhusaks	26
Väga lühikesed näidised inimestest ja biofilmidest	27
Lahendused	30
Biofilmid on väga mitmekesised	32
Näited sügava biofilmi mitmekesisuse kohta	32
Kahekordne löök: biokile ja ravimiresistentsusega bakterid	34
Antibiootikumiresistentsed "Superbakterid" on tõsine oht "väga lähitulevikus"	36
Lühike sõna Lyme'i biofilmide kohta	37
Lyme'i töbi (Borrelia) ja biofilmid	38
Kaks erinevat bakterit ühes biofilmis katastroof: tavaline Hambahügieen	39
Biofilmi hoolduse näidis	40
Bartonella ja Babesia biofilmid?	43
Puukide ja kirpude kaudu levivate biofilmide järelendus	43
Biofilmid ja kopsu- või põskkoopapõletikud: veski meditsiinikatastroofi anatoomia	45
Meeskonnasport: kui mitmest infektsioonist tekivad kõik biofilmid	46
Hoiatus: biofilmid ja lühikesed kiired meditsiinilised seansid	47

Eeterlikud õlid	48
Eugenoli põhitõed.....	52
Eugenool ja biofilmid	53
Linalool	57
Reserpiin	59
Biofilmi tapjate virnastamine	60
Terpenoidid	61
Allitsiin ja küüslauk	64
Serrapeptidaas	67
Lumbrokinaas	72
Nattokinaas	73
Terminalia chebula Retz	74
Vähk	77
Laktoferrin	82
Laktoferrini ostmine	84
Laktoferriniksülitooliga kombineeritud ravi	86
Traditsiooniliste suhkrute piiramine: põletiku ja biokilede langemine?	87
Suhkru ksülitool	88
Erütritool	90
Seleenorgaaniline	93
Kas magneesiumipuudus takistab biofilme?	94
Rasva piiramine biokilede raviks	97
Houttuynia cordata Thunb (HCT)	101
Biofilmi käivitavate kemikaalide proovid pöördusid tagasi bakterite vastu ..	104
Topeltravi	105
Taimsed allikad annavad harva ainult ühte kasulikku kemikaali	106
Nitroksoliin	107
Lüsosüüm	109

Aspiriin ja mittesteroidsed põletikuvastased ravimid	110
Asitromütsiin (Zithromax)	113
Hõbe	115
Gingerol	119
Stevia	121
Cumunda põhitoed	123
Cumanda ja biofilmid	125
Erütromütsiin	126
Kallis	128
Halb hingeõhk	134
Pööratud aminohapped õonestavad biofilme	136
Cathelicidin LL-37	139
RNAIII inhibeeriv ensüüm (RIP)	141
Tinidasool (Tindamax) ja Metronidasool (Flagyl)	142
Tuberkuloos (TB)	143
Uus tööriist tuberkuloosi tapmiseks?	144
Toidupulber, antibakteriaalne toime	145
Järeldus	146
Lisa	147
Dr Schalleri kliinilised väljaanded	153
Teised dr Schalleri raamatud	155
Vastutusest loobumine	169
Dr. Schalleriga ühenduse võtmine	170

Meditsiiniline revolutsioon

Biokileinfektsiooni teooria on sügav revolutsioon infektsioonide uurimisel, mis võivad olla valulikud, invaliidistavad ja tegelikult olenevalt vanusest kõige ohtlikumad.


Infektsioonid hakkavad meid tagasi viima aegadesse, mil inimesed surid lihtsatesse nakkustesse. Uus biokile nakatumise maailm võib tappa rohkem inimesi kui I ja II maailmasõda kokku, kui asjad ei muutu kiiresti nii arenenud kui ka arenemata riikides. Biokilede tähtsuse aeglase mõistmise ja seetõttu uute biokilelahenduste aeglase kasutuselevõtu tõttu arstide poolt võivad isegi tipparstid võtta biofilme tõsiselt alles siis, kui on tõestatud, et rohkem inimesi muutub puudega ja sureb nende tõttu. Praegu igatseb enamik biofilme kannatuste ja surma põhjusena. Seega on lahusteta biofilmid sama tõsised kui 19. sajandi lastehalvatus ilma vaktsiiniga ja ohvrite arvu poolest on need palju laastavamad kui HIV/AIDS.

Enamik baktereid elab kooslustes, millel on tavaliselt ainulaadsed kaitsvad biokiled. 1% inimesi nakatavatest või inimelu mõjutavatest bakteritest hõljub üksinda ja kui neid leitakse verest, ei leiaks neid koos ühegi biokile limaga.

National Institutes of Health hinnangul on enam kui 80% inimkehas esinevatest mikroobsetest infektsioonidest põhjustatud biokiledest, millest paljud põhjustavad kroonilisi ja korduvaid probleeme. Või on Glowackil õigus ja 99% bakteritest elab biofilmis? Olenemata sellest, kas kasutate hinnanguliselt NIH 80% või Glowacki 99%, on biofilmid infektsioonide puhul tõsiseltvõetavad.

Gyowacki R, Strek P, Zagórska-Swiezy K, Składziej J, Olej K, Hydzik-Sobocińska K, Miodoyski A. [Biofilm patsientidelt, kellel on krooniline rinosinusiit. Morfoloogilised SEM-uuringud].[Artikkel poolakeelne]. Otolaryngol Pol. 2008;62(3):305-10.

Sissejuhatavad biofilmipildid



Uus geneetiliselt ainulaadne biokilet tootv üherakuline parasiit nimega FL1953 või *Protomyxzoa rheumatica*. (See spetsiaalne määrdumine on parim viis nende üherakuliste parasiitide tuvastamiseks inimkehades, kuna DNA või PCR-test ei ole alati positiivne).


Selle pildi välisküljel olevad sada tumedat ovaali on 8 mikroni suurused punased verelibled (RBC). Keskmass on biokile pall, mille biokile massis on palju punaseid vereliblesid.

Seda ülaltoodud biokilet leidub tavaliselt inimestel, kellel on puukide kaudu levivad infektsioonid, nagu väga levinud *Bartonella*, Lyme'i tõve *Bor-relia* bakter ja surmav *Babesia*. Kuigi mõned puukide kaudu levivad haigused võivad olla teistest hullemad või levinumad kui teised, on kõik need potentsiaalselt surmavad, kui neid ei hävitata. See ülaltoodud parasiit on üherakuline infektsioon, mis on seotud *babeesia* ja malaariaga ning kui sellelt biokile eemaldatakse, näeb see välja nagu ebaküps malaaria. Haiguste tõrje keskuste andmetel on tegemist ainulaadse algloomaga. See ei ole *babeesia* ega malaaria. Seda infektsiooni nimetatakse FL1953 või *Protomyxzoa rheumatica*. See teeb tohutul hulgal biokilet ja sellel pildil olev tohtu keskmass sisaldab



Kuna me vaatleme erinevaid elundeid ja biokilede tekkepõhjuseid, ei tohiks me välja jätta biokileinfektsioonide vektorit, mida kannavad üle 200 elusolendi vähemalt kolmel kontinendil – iksoodide puuk. Sellel on vähemalt kaks tõsist biofilmitootjat: FL1953 ja väga keerulised geneetiliselt arenenud Lyme'i bakterid. Me alles õpime tundma kõiki võimalikke nakkusi, mida see kannab.

Pange tähele, et juuksed näevad välja nagu suur rohi, seega on see puuk sellest suurusest mürdosa. Kui kombineerite nähtamatuse hammustusega, millel on valuvaigisti, antihistamiin, antikoagulant ja põletikuvastane aine, on teil varjatud nakkuskandja. Üks puugi sülje kemikaal, Sialostatin L, on nii hea immuunsüsteemi pärssiv ensüüm, mis võib inhibeerida astmat (Horka 2012).



Koerad võivad olla inimese parimad sõbrad, kuid mitte siis, kui puudutate nende sülg ja mitte siis, kui nad toovad teie koju või autosse puuke või kirpe. Oletame, et igal väljaspool linna elaval koeral ja kassil on tõenäoliselt olnud puugi- või kirbuhammustus.



"Biofilmide" selgeks tegemine

Biokile on nagu peenraha oliiviõli basseini keskel ja õli välisservas on pipart, mis kujutab nakkust tapavaid rakke. Nad ei saa sisse kolida, et peenraha hävitada. Biokile bakterikooslused on enamiku inimeste nakkuste tavaline seisund. Meile on õpetatud, et nakkused on isoleeritud bakterid, mis hõljuvad ringi ja see on tõsine viga.

See näitab, kui kaugemale peame teaduses minema, kui bakterite peamine vorm biokile bakterite kogukonnad – on uus, kuid ülioluline kontseptsioon. Kui ma 2004. aastal koostas nimekirja kahekümne viiest võimalusest biofilmide hävitamiseks, ei olnud erilist huvi.

Tänapäeval on suutatus biofilme erinevate võimalustega hävitada sõna otseses mõttes tervisekatastroof.

Selle raamatu kirjutamise ja avaldamise eesmärk on luua taskukohane otsingupõhine valikute komplekt koos muude võimalike võimalustega, esitleda puhast lahenduste raamatut, mis pakub uusimaid võimalikke ja ajakohaseid lahendusi sadade seotud haiguste jaoks. biokiledega. Arstide, nakkusspetsialistide, naturopaatide, alternatiivmeditsiini koolide, eeterlike õlide praktiseerijate, nõelraviarstide, meditsiiniõdede või taimearstide tavapärase võimaluste korral võib bioloogilise kile barjääri olla täiesti võimatu eemaldada või läbi tungida.

Selle raamatuga loodame teenindada teid ja teie arsti/tervendajat, uurides praegu saadaolevaid võimalusi. Otsisime viimase viie aasta väljaandeid PubMedist - tohutust arstiteaduse andmebaasist - "biofilmi töötlemiseks". Valikute valik on muljetavaldav ja mitte alati midagi, mida võite oodata. Selle raamatu eesmärk on anda teile laialdased võimalused teie kannatuste, puude ja isegi surma ärahoidmiseks.

Pärast aastaid kestnud uurimistööd olen aru saanud, et biofilmi nakkushaiguste "eksperdid" võivad olla sõja ammu kaotanud ja tegelikult ei pruugi paljud kõigist lahingutest kunagi teadlikud olla. Pa-

Väga lühikesed näidised inimestest ja biofilmidest

2004. aastal paranes Richard Longland pärast lüüsisambaoperatsiooni müstilisest haigusest väga halvasti. Järgnevatel kuudel kannatas ta paljude probleemide all – peavalud, liigesevalu ning hiljem südame- ja ajuprobleemid, jõhker väsimus ja mõtlemisraskused.

Meditsiinisüsteem oli talle vastu, kuid lõpuks, 2007. aastal, raviti teda mükoplasmaga, mis tekkis võimalikust operatsioonist, mis tahes kohast haiglas või avalikus kohas või puugist.

Enamik minu patsientidest on enne minu juurde tulekut käinud 3–200 arsti juures. Ma saan tema kogemusest aru. Hr Longland pidi diagnoosi saamiseks pöörduma üle kahekümne arsti poole. Sellel raskel perioodil lõi ta suurepärase filmi "Miks ma olen nii haige?" Ta on patsientide tšempion farmatseutiliste ja naturopaatiliste ainete kasutamises, et vabastada oma keha süsteemsetest bakteriaalsetest biofilmidest.

Edward on 78-aastane ja tal on kolm tüdruku ja kaheksa lapselast. Ta viidi õhupuuduse tõttu haiglasse. Tal on halb kopsupõletik või infektsioon kopsus. Ta läheb hullemaks. Üksikisikud on paranenud, kasutades aineid, mis võivad paljusid biokilega kaitstud kopsupõletikke.

Linda on juba mõned aastad väsinud ja kooliga hätta jäänud. Hiljuti avastasin, et tal on mitu puugiinfektsiooni, mis on põhjustanud üle viieteistkümne laboritulemuse ebanormaalseks. Eile ta helistas ja põlvetaguse valu tõttu käskisin tal kiirabisse minna. Vähem kui ööpäevaga avastati tal kopsudes ja jalgades 23 trombi. Ta kahtlustab, et see on babeesia, põletik ja FL1953. Meil olid agendid, kes tapsid need agendid, sealhulgas FL1953, 2006. aastal.

It would be an error to say that nattokinase, lumbrokinase, serrapeptidase, EDTA, gentamicin, vancomycin, Samento, Banderol, olive products, poorly known herbs with fair lab testing in humans, clove bud oil, diet, chelation, three to four part amino acid mixes, NAC, Rife, diet changes or a vast range of other options not listed, will **work for all biofilms**. For example, an elderly patient dying of a lung infection or another person with painful and treatment-resistant sinus infection *will not* have the same biofilm.

As a trend, trying different options to destroy a biofilm is less dangerous than allowing it to spread.

A Brief Word on Biofilms in Lyme

At times, individuals who have tick- and flea-borne infections, like Bartonella, Babesia and Borrelia (Lyme disease), can feel their treatment is minimal or incomplete. Debates rage over the diagnosis and treatment of Lyme and tick-borne diseases; whether the pain is from residual dead infection incorporated into tissue or one of the many infections carried by the I. scapularis tick, we still have patients' misery.

After writing **twelve books** which include many pages on non-Borrelia infections, “Lyme testing” seems like alphabet testing in which ***one only looks for the vowel “a.”*** Due to the lack of acceptance of the number and complexity of tick-borne infections, there is a lack of up to date education, leaving quality medical doctors to evaluate tick and flea infections in the ***abstract***, by which I mean that they very falsely and sadly do not realize the full magnitude of ***“the alphabet.”***

Specifically, they “diagnose” by ignoring inflammation alterations, nutrient changes, hormone deficits, immunity changes caused by tick-borne infections, and chemicals made or suppressed by direct tick and flea infectious agents. I discuss these in my three most recent tick and flea infection books. All are available in English. All can be found free through inter-library loan, for less than \$20 USD, or at www.personal-consult.com under the “free books” button. No one can expect to become an expert in this massive area after reading any guide or merely going to ten conferences, because these cluster infections impact twenty areas of medical and scientific knowledge.

In the last four years, researchers like **Dr. Eva Sapi have shown Lyme is like some other spirochetes—it has biofilms. These are very tough biofilms to defeat unless caught in the “acute stage.”** A tough, “mature biofilm” allows organisms to **“laugh at” many antibiotics.**

Some medical professionals interested in Lyme often ignore the immune suppressing Bartonella bacterium, which is more common than Lyme. Ignoring coinfections may increase the risk of fatality with Babesia and possibly **FL1953**. These healers also may not realize that the highly

genetically complex Lyme spirochete appears to have a troublesome biofilm. Performing a simple direct test at laboratory companies whose testing kits have reduced sensitivity will probably result in more negatives for tick-borne diseases. The ultimate result is anti-science and anti-truth. Searching for tick infections with one test is like writing in “Lincoln” at the next presidential election.

Lyme Disease (*Borrelia*) and Biofilms

Several researchers believe *Borrelia burgdorferi*, the active agent of Lyme disease, has biofilms. Lyme organism biofilms have been found in culture and in the tick gut. Lyme cysts and biofilms have also been noted in patient skin biopsies using focus floating microscopy according to Dr. Eisendle publishing in the *American Journal of Pathology*.

Further, we see in Lyme that biofilm formation is dependent on cyclic di-GMP expression and we see that in Lyme (Stricker and Johnson).

Brihuega B, Samartino L, Auteri C, Venzano A, Caimi K. In vivo cell aggregations of a recent swine biofilm-forming isolate of *Leptospira interrogans* strain from Argentina. *Rev Argent Microbiol*. 2012 Jul-Sep;44(3):138-43. PMID:23102459

Cogoni V, Morgan-Smith A, Fenno JC, Jenkinson HF, Dymock D. *Treponema denticola* chymotrypsin-like proteinase (CTLP) integrates spirochaetes within oral microbial communities. *Microbiology*. 2012 Mar;158(Pt 3):759-70. Epub 2012 Feb 7. PMID:22313692

Sapi E, Kaur N, Anyanwu S, Luecke DF, Datar A, Patel S, Rossi M, Stricker RB. Evaluation of in-vitro antibiotic susceptibility of different morphological forms of *Borrelia burgdorferi*. *Infect Drug Resist*. 2011;4:97-113. Epub 2011 May 3. PMID:21753890

Stricker RB, Johnson L. Lyme disease: the next decade. *Infect Drug resist*. 2011; 4: 1-9. PMID: 21694904

Sapi E, Bastian SL, Mpoy CM, Scott S, Rattelle A, Pabbati N, Poruri A, Burugu D, Theophilus PA, Pham TV, Datar A, Dhaliwal NK, MacDonald A, Rossi MJ, Sinha SK, Luecke DF. Characterization of biofilm formation by *Borrelia burgdorferi* in vitro. *PLoS One*. 2012;7(10):e48277. Epub 2012 Oct 24. PMID:23110225

lease of bacteria in the human body will be like a dangerous tornado in a field. It is a wise concern.

For these two problems regarding biofilm-held infections suddenly being released, here are useful solutions:

1. You need many infection killing options for use since more is better to prevent “seeding” of dispersed infection.
2. You want the biofilm killing options to destroy biofilms by different mechanisms. This makes the dispersed seeded infections naked to the immune system.
3. Biofilm tools are given initially at low doses and then increased gradually to large doses since often in the beginning patients have massive inflammation and a drastic increase in killing of biofilm organisms in a short time could cause trouble with bone marrow, liver, heart, eye, or kidney issues, or merely create more dead infectious debris resulting in patient misery.
4. You may need to pulse (use every other day) or fully stop this treatment because once a wave of biofilm eroding agents strips off or severely damages a biofilm of an infection, the same antibiotics that were useless in the past can become very effective.
5. There is no single master biofilm destroyer, yet some are broader than others.

Bartonella and Babesia Biofilms?

Most people have heard of the profoundly common tick infection Lyme disease, but they may not know Bartonella is more common than Lyme and is carried by far more vectors (Breitschwerdt). Babesia decimated the cattle population in the southern United States many decades ago and is more dangerous in humans than Lyme.

Currently, we have no solid data showing Bartonella and Babesia have biofilms.


Tick and Flea-Borne Biofilms Conclusion

Below you will see that mouth spirochetes routinely have biofilms. Another spirochete is Leptospira which is able to make biofilms in many environments and may contribute to lost pregnancy in mammals (Brihuega).

In terms of tick and flea infection biofilms, I would focus on **FL1953** (Protomyxzoa) and Lyme, since both have been known and treated by us since 2006, though the former was killed without knowing its genetic uniqueness. We are learning what decreases their biofilm pathology and have agents that should work if one is open to look at diverse approaches. A synthetic “antibiotic only approach” to biofilms, including antibiotics targeted to hit biofilms, might be similar to typing with one finger.

There are herbalists, such as Stephen Buhner, who propose selected herbs to treat some tick infections. And, in terms of **primary treating herbs to kill organisms**, there are also credible options that are not always herbal in use for a tick or flea infection. We will continue to use **advanced lab testing**, typically only allowed under physician supervision, to determine by serious extensive **indirect blood exam** biochemistry tests to see which infection is actually destroyed in people experiencing benefit from herbal therapy. In any event, I enjoyed this line from Buhner: *“I can’t really say what will clear all biofilms.”*

Veel üks biofilmi näidispilt



Tume IOVId oVlls ai:e punane veri "°Us (-blade upp« vares), Tho "tehe" diat algab llo paremalt madalalt« 'o'11«, niida vasaku ütemise oomeri poole, haige biofilil, \ rnlffffial IOV\ll' niidab— poi111i11g väikesele bam:riumile. (Piy Labaratoriea)

Eugenoli põhitõed

Eugenooli leidub paljudes eeterlikes õlides ja maitsetaimedes. Näiteks leidub seda tugevalt nelgipungade eeterlikus õlis, kuid väiksemas annuses kaneelilehtedes ja selle eeterlikus õlis. PubChemi andmetel leidub seda ka pimento-, bay-, sassafras-, massoy-kooreõlis, kampriõlis ja chamchwi taimedes. Tugevus ja kontsentratsioon varieeruvad olenevalt allikast ja ekstraheerimismeetodist suuresti. Lisaks ei ole see lihtsalt võimas biofilm aine; sellel on muid hämmastavaid omadusi, nagu viirusevastane toime ja väh



Näiteks näitasid Tragoolpua ja Jatisatienr, et eugenool mõjutab suu- ja suguelundite herpest sõltuvalt liigist, tüvest ja muudest teguritest. Nad tegid selgeks, et eeterlik õli võib olla võimsam kui lihtne ekstrakt. Tõepoolest, suu- ja suguelundite herpes, vastavalt HSV-1 ja HSV-2, ei saanud euge-nol juuresolekul paljuneda. Al-Sharif on näidanud märkimisväärset vähi mõju. Väga madalal kontsentratsioonil (2 μ M) on spetsiifiline toksilisus erinevate rinnavähirakkude suhtes. Seda tapvat toimet vahendas vähirakkude surmatee esilekutsumine ning E2F 1 ja surviviini taseme langus – kaks molekuli, mis on rakkude ellujäämiseks olulised. See takistas k

genes. Importantly, these anti-proliferative and pro-cancer cell death effects were also observed inside body grafts placed in non-human animals.

<http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=3314>

Tragoalpua Y, Jatisatiennr A. Anti-herpes simplex virus activities of *Eugenia caryophyllus* (Spreng.) Bullock & S. G. Harrison and essential oil, eugenol. *Phytother Res.* 2007; 21(12):1153-8.

Al-Sharif I, Remmal A, Aboussekhra A. Eugenol triggers apoptosis in breast cancer cells through E2F1/survivin down-regulation. *BMC Cancer.* 2013 Dec 13;13(1):600. [Epub ahead of print]

Eugenol and Biofilms

Recently, Dr. Zhou has reminded us of a special process that is involved in the formation of dangerous biofilms. Basically, many bacteria have a “chatty” way of talking to other cells such as other bacteria. So, bacteria use chemicals or cause other bacteria to make chemicals to help them survive and often act to harm you or a loved one.

Eugenol is so effective that at very low amounts, it still disrupted bacteria chemical communication. This is very important in a biofilm destroying agent. If cells cannot communicate, it is doubtful they can form communities. Biofilms are community creations. **Further, eugenol at very low doses, called “sub-inhibitory concentrations” inhibited biofilm formation.**

One type of biofilm research being conducted compares biofilm killers head to head. The results are not always the same, perhaps in part because the infections are not always the same. Note that in an Epub abstract before publication, Malic explains that the best essential oil for urinary catheters, with or without biofilms, against fourteen different bacteria was eugenol. This is why I believe this substance is a “double killer.” It can defeat many biofilms, and then kill the organism making the biofilm. Finally, in this study, eugenol did better than tea tree oil.

Linalool

According to the Merriam-Webster dictionary, the word linalool is derived from a Medieval Latin phrase meaning “wood of the aloe.” Linalool has a nice smelling alcohol and essential oils. It is used in perfumes, soaps, and flavoring materials.

In terms of biofilms, it seems to be most effective when **the essential oil part** is used, which has **the most evidence of killing Candida albicans**. (*Candida albicans* is the cause of yeast infections.) Yet, again, it is the essential oil fraction that not only **inhibits the growth** of *Candida albicans* but also of the bacteria *Lactobacillus casei*, *Staphylococcus aureus*, *Streptococcus sobrinus*, *Porphyromonas gingivalis* and *Streptococcus mutans* cell suspensions, all of them associated with oral cavity disease, according to Alviano and Mendonça-Filho. Yet, Budzyńska reported this essential oil did not fully remove biofilms formed by *Staphylococcus aureus* (ATCC 29213) and *Escherichia coli* (NCTC 8196) on the surface of routine medical materials such as urinary catheters, infusion tubes and surgical mesh.

Hsu found that linalool could be effective against *Candida albicans* due to its many genetic blocking effects. For example, using a scanning electron microscope and other technology, many signs of the effect of linalool to destroy *Candida* or inhibit its growth could be noted. Hsu found blocking actions against genes involving adhesion production and the formation of “branches” or the mold’s hyphae were both decreased by linalool.

<http://www.merriam-webster.com/dictionary/linalool>

Budzyńska A, Wieckowska-Szakiel M, Sadowska B, Kalemba D, Różalska B. Antibiofilm activity of selected plant essential oils and their major components. *Pol J Microbiol*. 2011;60(1):35-41. PMID:21630572

Alviano WS, Mendonça-Filho RR, Alviano DS, Bizzo HR, Souto-Pradón T, Rodrigues ML, Bolognese AM, Alviano CS, Souza MM. Antimicrobial activity of *Croton cajucara* Benth linalool-rich essential oil on artificial biofilms and planktonic microorganisms. *Oral Microbiol Immunol*. 2005 Apr;20(2):101-5.

Reserpine

Reserpine is a substance found in the roots of some types of Rauwolfia that has been made into a traditional medicine. It is used to lower high blood pressure and help with psychotic symptoms, but side effects have limited its use.

While it may not be comfortable to use at modest or high dosing, very low dosing, according to Magesh, showed it to be profoundly powerful against *Klebsiella pneumoniae*. In one report, he used reserpine and was able to stop biofilms in this pneumoniae infection at a fraction of the dose thought to inhibit growth.

Specifically, a tiny fraction of this drug, a mere 0.0156 mg/ml, stopped biofilm production in *Klebsiella pneumoniae*. So, it may be possible that we have another example of a medical truth I use every day:

“Change the dose and you change the drug or herb.”

In this case, perhaps it is possible that 1/10th of the lowest size tablet, 0.1 mg, could harm *Klebsiella* and other infections and be safe for the patient. However, the raw materials for making it may be hard to find some months according to ASHP who tracks pharmacy shortages.

Magesh H, Kumar A, Alam A, Priyam, Sekar U, Sumantran VN, Vaidyanathan R. Identification of natural compounds which inhibit biofilm formation in clinical isolates of *Klebsiella pneumoniae*. *Indian J Exp Biol*. 2013 Sep;51(9):764-72.

<http://www.ashp.org/DrugShortages/Current/Bulletin.aspx?id=975>

“Stacking” Biofilm Killers

While physicians may ponder the problems caused by biofilms in practice, I rarely encounter the doctor who understands that it is usually better to have more than one treatment. In the article below, **oral bio-film infections were controlled best by three agents, not merely one.** For example, Alves explains that when you are going to irrigate or clean a root canal area, that two mouth bacteria infections protected by their biofilms have these same film barriers decreased significantly by treatment with farnesol, xylitol and lactoferrin together.

The same results were found in wounds. One of the best treatments for wounds is the use of a silver-based wound dressing or bandage, together with a gel containing xylitol and lactoferrin (Ammons).

Alves FR, Silva MG, Rôças IN, Siqueira JF Jr. Biofilm biomass disruption by natural substances with potential for endodontic use. *Braz Oral Res.* 2013 Jan-Feb;27(1):20-5. PMID:23306623

Ammons MC, Ward LS, James GA. Anti-biofilm efficacy of a lactoferrin/xylitol wound hydrogel used in combination with silver wound dressings. *Int Wound J.* 2011 Jun;8(3):268-73. Epub 2011 Apr 1. PMID:21457463

Terpenoids

I would like to mention a class of options that come from a familiar substance, chemicals from tea tree oil. We have already mentioned linalool which is part of this class individually, since it comes up as a leading biofilm killer. According to Raut, as many as 14 terpenoids derived from tea tree oil inhibit biofilms, and α -terpineol, nerol, isopulegol, carvone, linalool, α -thujone and farnesol are worthy of special note. Eight terpenoids have effects on **mature** yeast biofilms (*Candida albicans*).

A study by Ramage shows tea tree oil (TTO), terpinen-4-ol (T-4-ol), and α -terpineol displaying potent activity against 69 biofilm-forming *Candida* strains, of which T-4-ol and α -terpineol displayed rapid kill action.

Of these three, T-4-ol displayed no significant toxicity to cells. These data provide further laboratory evidence that TTO and its derivative components, specifically T-4-ol, exhibit strong antimicrobial properties against fungal biofilms. Further, T-4-ol appears to possess safety advantages over the complete essential oil (TTO) and may be suitable for prevention and treatment of established oral and upper throat cavity candidosis. Certain terpenoids are components of spices or food ingredients generally regarded as safe (GRAS) (Pauli 2006).

In another study, several chemicals from plants were tried against two very common bacteria (Budzyńska), *Staphylococcus aureus* (ATCC 29213) and *Escherichia coli* (NCTC 8196), both with biofilms on the surface of **routine** medical products, i.e., urinary catheter, infusion tube and surgical mesh. All three are present in most advanced hospitals and other settings. Surgical mesh was the surface most prone to persistent colonization since the biofilms that formed on it, both by *S. aureus* and *E. coli*, were difficult to destroy.

Melaleuca alternifolia is the source of Tea Tree Oil (TTO). *Lavandula angustifolia* yields Lavender, English Lavender and True Lavender (LEO). *Melissa officinalis* is Lemon balm (MEO). Tea Tree oil, Lemon balm, α -terpineol and terpinen-4-ol showed stronger anti-biofilm

Allicin and Garlic

Garlic has been used as a medicine throughout human history. Allicin is considered one of the medically useful components of garlic. Other useful components are discussed in Chinese language pharmacology texts.

As early as 2003, the use of allicin against *Staphylococcus epidermidis* had reported effects on biofilm formation at low dosing. Pérez-Giraldo reported that lab testing showed that allicin diminished biofilm formations.

Lihua reported ten years later that allicin impacts *Pseudomonas aeruginosa* biofilm. This is hardly casual information, since *P. aeruginosa* is likely resistant to multiple antibiotics, and this resistance may be due to biofilms. Organosulfur allicin has been shown to inhibit surface-adherence of bacteria and Lihua demonstrated that allicin could inhibit early bacterial adhesion which is a first step to bacterial community formation, usually just before biofilm production.

Other researchers isolated various components of garlic and tested the most active components. The following three components were examined:

1. garlic extract
2. allicin
3. diallyl sulfide (DAS)

They were tested against the serious mouth and dental infection *Aggregatibacter actinomycetemcomitans*, the primary cause of severe aggressive periodontitis and other non-oral infections.

Lumbrokinase

We appreciate that some people interested in progressive medicine feel this enzyme, Lumbrokinase, is a useful substance. Some have suggested it is useful in the removal of biofilms. If that is true, we had trouble finding the evidence for that position. However, it does seem that some researchers see a potential for this enzyme to “digest” pathological clots. This possibility seems to have some support, and at this time we will only wait for further research. Since we are only proposing biofilm options that are supported by research and since human use is just starting in research settings, we do not promote this agent at this time.

Ryu GH, Park S, Han DK, Kim YH, Min B. Antithrombotic activity of a lumbrokinase immobilized polyurethane surface. *ASAIO J.* 1993 Jul-Sep;39(3):M314-8. PMID:8268550

Kim JS, Kang JK, Chang HC, Lee M, Kim GS, Lee DK, Kim ST, Kim M, Park S. The thrombolytic effect of lumbrokinase is not as potent as urokinase in a rabbit cerebral embolism model. *J Korean Med Sci.* 1993 Apr;8(2):117-20. PMID: 8397927

Mihara H, Sumi H, Yoneta T, Mizumoto H, Ikeda R, Seiki M, Maruyama M. A novel fibrinolytic enzyme extracted from the earthworm, *Lumbricus rubellus*. *Jpn J Physiol.* 1991;41(3):461-72. PMID:1960890

Wang KY, Tull L, Cooper E, Wang N, Liu D. Recombinant Protein Production of Earthworm Lumbrokinase for Potential Antithrombotic Application. *Evid Based Complement Alternat Med.* 2013;2013:783971. Epub 2013 Dec 12. Review. PMID:24416067

Cao YJ, Zhang X, Wang WH, Zhai WQ, Qian JF, Wang JS, Chen J, You NX, Zhao Z, Wu QY, Xu Y, Yuan L, Li RX, Liu CF. Oral fibrinogen-depleting agent lumbrokinase for secondary ischemic stroke prevention: results from a multicenter, randomized, parallel-group and controlled clinical trial. *Chin Med J (Engl).* 2013 Nov;126(21):4060-5. PMID:24229674

Huang CY, Kuo WW, Liao HE, Lin YM, Kuo CH, Tsai FJ, Tsai CH, Chen JL, Lin JY. Correction to Lumbrokinase Attenuates Side-Stream-Smoke-Induced Apoptosis and Autophagy in Young Hamster Hippocampus: Correlated with eNOS Induction and NF κ B/iNOS/COX-2 Signaling Suppression. *Chem Res Toxicol.* 2013 Jul 15;26(7):1126. Epub 2013 Jun 7. PMID:23746067

tract also made the pneumonia far more susceptible to the antibiotic tobramycin. Further, genes involved with resistance to antibiotics were down-regulated.

- Bag published that highly resistant urine organ infections were more vulnerable to treatment with *T. chebula* but proposed this is due to its ability to collect iron, since adding iron reduced its effect. However, Bag only tested one of many chemicals from this fruit, and I would suggest other components may have antibacterial action and work by other means.
- Four carefully chosen antibacterial plants (*P. guajava*, *T. chebula*, *A. aspera*, and *M. elengi*) are combined with four solvent extracts (hexane, ethyl acetate, ethanol, and methanol) by Kamal Rai Aneja, who initially evaluated their anti-cavity activity against *S. mutans*. All four of the plants showed activity against *S. mutans*. Ethyl acetate extracts of the four plants showed high antibacterial activity against *S. mutans*, superior to the other solvent extracts. Further, *T. chebula* ethyl acetate extract acts as an effective anti-cavity agent by inhibiting *S. mutans* and *C. albicans*. However, we were unable to find evidence if the benefit of these chemicals involved biofilm removal.

In conclusion, we appreciate that this medicine is proposed to both dissolve Lyme biofilms and also destroy the underlying Lyme bacteria. We offer no opinion on this belief. We do not want to oppose or support its use in terms of biofilm ability. It appears this fruit does act on the bacteria biofilm of *P. aeruginosa*, but Lyme bacteria are not the same as *P. aeruginosa* bacteria. Lyme is also profoundly more genetically complex than a “relative” spirochete bacterium, syphilis.

Therefore, while we do note that this medicine has antibacterial and cell protection actions, and **we accept some patients feel better**, we presently cannot say it is due to biofilm removal in those with tick-borne infections.

Cancer

Cancer has many causes. Some things increase your risk and other things can decrease your risk. It is rarely pure genetics, even in those with genetic vulnerability. We know some types of plastics increase rates of breast cancer. We know the 200 poisons in cigarettes cause lung cancer. We know various chemicals made by various companies can increase cancer, despite the reality that most US and international chemicals have limited or no top research on their safety.

I like my dental hygienist. And, I like making sure my gums and teeth are “safe.” Why? At first it was because I want to have teeth in twenty years. But, she correctly reminds me that heart attacks are increased by gum disease which is routine in many countries.

Yet, even this passionate healer was not aware of the role of biofilms in cancer. Yes, I said cancer. We are only beginning to understand the role of infections in triggering cancer diseases.

Many years ago, I was working with a physician who asked me to help research possible cures for his cancer. Eventually, that cure was found and written up, taking over 200 hours and many months to complete, with the help of a top medical editor in North America—the former editor of the *Journal of the American Medical Society* and forty other journals, specifically, George Lundberg, who worked feverishly to get this death disorder cure in print ASAP (Schaller).

Years later, he asked me to write a follow up, and we had found that over eight top infection specialists in the United States had missed Babesia, a common parasite that is harder to kill than malaria and which can occasionally increase eosinophils (Schaller). The patient’s trouble included the fact that he had so many eosinophils, his blood could clot quickly. The point? Eosinophils are a type of white blood cell designed to kill parasites. The man’s disorder (HES) Idiopathic Hypereosinophilic Syndrome, which is often fatal and means that eosinophils reproduce out of control, was primed by a Babesia infection. Not all patients with HES also have a Babesia infection, but after writing six books which

Lactoferrin Xylitol Combination Treatment

In a fascinating look at this proposed double treatment, Mary Ammons shares that treatment of *Pseudomonas aeruginosa* biofilm with both lactoferrin and xylitol inhibits the ability of bacteria to respond to damage resulting from lactoferrin iron chelation.

Pseudomonas aeruginosa has been identified as the most common biofilm-forming infection in chronic wounds. The immune stimulating molecule lactoferrin and the rare sugar alcohol xylitol, together, were effective in the lab against *P. aeruginosa* biofilms.

How? Lactoferrin iron chelation was identified as the primary means by which lactoferrin undermines the bacterial membrane. Amazingly, this combination showed huge alterations in the expression of the bacteria's genes, but these changes are too complex for a summary. The findings mean that critical chemicals made by *P. aeruginosa* had changed.

Siderophore detection verified that xylitol is the component of this unique double treatment that inhibits the ability of the bacteria to produce siderophores under conditions of iron restriction. Siderophores sound complicated—here is the simple meaning: they are some of the strongest iron binders in the world and they are made by bacteria, viruses and fungi.

The study concludes with two points:

1. Lactoferrin treatment of *P. aeruginosa* biofilms results in destabilization of the bacterial cell membrane through iron chelation.
2. Combining lactoferrin and xylitol inhibits the ability of *P. aeruginosa* biofilms to respond to environmental iron restriction.

Access to iron is profoundly hard for bacteria when this combination is used.

Erythritol

Erythritol is an amazing sugar. For example, when it was given to children head-to-head with xylitol or sorbitol it was clearly superior. Here is a summary of the research:

Runnel writes: “Three-year consumption of erythritol-containing candies by initially 7- to 8-year old children was associated with reduced plaque growth, lower levels of plaque acetic acid and propionic acid, and reduced oral counts of mutans streptococci compared with the consumption of xylitol or sorbitol candies.”

In a similar way, Japanese researchers show highly advanced reasons for erythritol superiority over xylitol and sorbitol (Hashino). While this study is very dense, let me at least try to list the stunning findings:

1. By advanced confocal microscopic observations, the most effective sugar used to reduce *P. gingivalis* accumulation onto an *S. gordonii* substratum was erythritol, as compared with xylitol and sorbitol.
2. In addition, erythritol moderately suppressed *S. gordonii* monotypic biofilm formation.
3. To examine the inhibitory effects of erythritol, they analyzed the metabolomic profiles of erythritol-treated *P. gingivalis* and *S. gordonii* cells. Metabolome analyses showed that a number of critical bacteria chemicals were decreased by erythritol.
4. Next, metabolites of erythritol- and sorbitol-treated cells were examined. Erythritol significantly decreased the levels of *P. gingivalis* dipeptides. They tended to be increased by sorbitol.

Amazingly, it appears erythritol has inhibitory effects on two diverse species with biofilms, and it acts by at least five very distinct mechanisms.

Dowd reported that biofilm formation was completely inhibited in a standard wound approach by 10% erythritol in either of the two San-

Does Magnesium Deprivation Hinder Biofilms?

Before we decide to remove an element that is used in vast numbers of important enzymes, we have to have a foundation. First, in some basic physiology texts, calcium displaces magnesium inside human cells. My impression of this research is that suboptimal magnesium increases systemic inflammation, vascular death such as heart attacks, and cancer. Dibaba shows that the higher the magnesium in diet the lower C-reactive protein. This protein is associated with inflammation. If you lower inflammation you decrease deaths.

Qu pooled studies of approximately a half a million people to examine the results. The greatest risk reduction occurred when magnesium intake increased from 150 to 400 mg/day. A significant inverse association was found between dietary magnesium intake and total cardiovascular events. Serum magnesium concentrations are linearly and inversely associated with the risk of cardiovascular troubles such as heart attacks and brain strokes. Since magnesium is poorly absorbed even when chelated to an amino acid, it is perhaps useful to note the useful dose was 400 mg, when compared to minimal benefit from 150 mg orally.

Del Gobbo also examined vast studies and wrote: “Clinical hypomagnesemia and experimental restriction of dietary magnesium increase cardiac arrhythmias.” Deadly ischemic heart disease, in which a person dies due to poorly oxygenated blood reaching the entire heart, was more common in those with no magnesium supplementation or very low oral magnesium dosing. Simply, “circulating and dietary magnesium are inversely associated with [cardiovascular disease].” Further, Qu shows, in another study, a significant drop in intestinal cancers with a reasonable magnesium intake. While we may not know the mechanism for these useful findings, they are not felt to be due to chance.

Song and Leff clearly show why a small number of scientists and physicians have pondered lowering human magnesium Mg^{2+} levels. They remind us that Mg^{2+} can influence bacterial adhesion, which is part of biofilm process. In their study, the bacterium *Pseudomonas fluorescens* was used to investigate the influence of Mg^{2+} on biofilm growth.

Nitroxoline

We are not going to spend significant time on this fifty year-old antibiotic because it is not used in many countries, and it is a quinolone, and quinolones all seem to have serious risk of tendon damage. For example, it is possible nitroxoline has the same risks as other quinolones (www.drugbank.ca/drugs/DB01422).

Quinolones easily enter cells and are often used to treat intracellular pathogens such as *Mycoplasma pneumoniae*.

The FDA has increased warnings regarding side effects since the drugs were first approved. I just want to focus on three side effects that might not be routine but are possible risks with many quinolones:

- **Damage to nerves outside the brain:** This could present as sensory nerve or muscle nerve injury causing paresthesias, hypoaesthesias, dysesthesias, and weakness. New pain, burning, tingling, numbness and/or weakness, or new decreased abilities to detect light touch, pain, temperature, position sense, vibratory sensation, and/or motor strength are basic nerve functions and show damage; these are reasons to stop taking the drug.
- **Tendon damage:** While some focus on the Achilles tendon, actual tears of tendons have occurred in the hand, the shoulder, the thigh, or other locations. Some are helped with surgery. Other patients feel the surgical or other treatment still leaves them with damage. It is believed by some that the use of prednisone and other cortical steroids meant to drop inflammation increases the risk of tendon damage. Perhaps this is especially true in older seniors. Surprisingly, tendons can rupture after the medication is stopped. Some have suggested that IV, transdermal or sublingual magnesium might decrease the risk, but I am not aware this hypothesis has been proven (Schaller).

Aspirin and NSAIDS

We have previously said it is best to see biofilms like a key, and using AIDS as an example, it was only after AZT in 1996 with **the arrival of protease inhibitors that those quickly dying, experienced a “Lazarus effect,” in which AIDS patients who looked to be ready to die recovered markedly in 30 days.** Medications used in AIDS are tough medications, even if they are miracles. Some may question offering a section on the tough medications aspirin and NSAIDS.

While we appreciate that aspirin and various other over the counter NSAIDS may not be optimal, perhaps due to concerns of liver, kidney or ulcer issues, we are discussing infections that invade and cannot be stopped by your body. You might need all the help you can get. So we offer some synthetic options here that may offer help against a top killing and disabling problem—**biofilm-protected** infections.

For example, fluconazole-resistant *Candida* is increasing worldwide. Fluconazole is also called Diflucan. Biofilms are one reason for a decreased effect in treatment. Aspirin, diclofenac, ketoprofen, tenoxicam, and ketorolac all undermined biofilms or their processes. They all reduced fungal adhesion, and increased biofilm detachment with low concentrations of anti-inflammatory agents. Microscopic examination confirmed the tested drugs had a significant effect on reduction of *Candida* adhesion and biofilm development. The drugs also made fluconazole work more effectively against fluconazole-resistant *C. albicans* (Abdelmegeed).

Another useful way to involve aspirin is by teaming it up with the chelation chemical EDTA. Both aspirin and EDTA possess broad antimicrobial activity for biofilm cultures. Aspirin used for 24 hours was successful in eradicating *P. aeruginosa*, *E. coli* and *C. albicans* biofilms. Moreover, exposure to the Aspirin-EDTA combination completely destroyed bacterial biofilms after only four hours in simulation lab testing (Al-Bakri).

Azithromycin (Zithromax)

This medication is almost a household name and is known as the “Z-Pak” which contains brand name Zithromax pills that are still in use today. Despite being in use many years and used very routinely, this medication still has a strong use in addressing biofilms.

For example, Maezono showed that azithromycin was markedly superior compared to other routine antibiotics in killing gum infection bacteria. Specifically, azithromycin at **very low dosing** undermined four strains of *Porphyromonas gingivalis*. This determination involved the use of two fascinating techniques.

Azithromycin dropped the bacteria “gasoline” or ATP in the bacteria, which means the bacteria had decreased function or were dead. Cyanide kills humans in part due to dropping ATP levels—it is not a trivial substance. Further, the power of azithromycin was seen clearly with a confocal laser scanning microscope, which has the ability that the long name suggests—seeing the decreased amount of bacteria.

One of the most common hospital infection risks is MRSA; it causes a number of potentially deadly diseases. This “MRSA” simply means routine staph aureus is no longer able to be killed or it is resistant to methicillin, so it reproduces unchecked. Azithromycin is proposed as one solution to MRSA based partly on its biofilm defeating abilities at very low dosing.


Gui shows that azithromycin was active against methicillin-resistant *Staphylococcus aureus* (MRSA) strains. It reduced the production of α -hemolysin and biofilm formation at very low “sub-inhibitory” concentrations. So, azithromycin may be useful in the treatment of α -hemolysin-producing and biofilm-forming MRSA infections.

Maezono H, Noiri Y, Asahi Y, Yamaguchi M, Yamamoto R, Izutani N, Azakami H, Ebisu S. Antibiofilm effects of azithromycin and erythromycin on *Porphyromonas gingivalis*. *Antimicrob Agents Chemother*. 2011 Dec;55(12):5887-92. Epub 2011 Sep 12. PMID:21911560

Hõbedane

Haavade biokilede vastu kasutatav hõbedatöötlus on olnud selgelt tõhus. Tõepoolest, 1% hõbedast kreemi on kogu maailmas edukalt kasutatud põnnihaigete infektsioonide raviks ja ennetamiseks.

Rahvusvahelise haavainfektsiooni instituudi ülevaade näitab, et andmed viitavad endiselt hõbedale kui parimale raviviisile. Näiteks katsetas Monteiro kolloidhõbedat seente biokilede vastu. Selle töö järelendus on väga kindel: olenemata uuringus kasutatud kontsentratsioonidest mõjutas hõbe Candida biokilede matriksi koostist ja struktuuri.



3-mõõtmeline renderdatud lähivõte Candida albicansist.

Cumanda and Biofilms

Dr. Eva Sapi and her colleagues found in their superior laboratory that cumanda had some mild killing effects on the Lyme bacteria, but more importantly for this book, Lyme **biofilm** communities grown in her lab were reduced 43% by this herb at low dosing. The dosing for a dynamic human or animal body was not explored or proposed by this researcher or any other researcher as of February 2014. Searching by its Latin and popular name did not yield any articles relevant for use on infections.

Finally, while Lyme disease is a common and disabling infection, it is hardly the only infectious agent in the many infections carried by Ixodes ticks. While this preliminary research is very useful, it is possible cumanda may have impact inside a body for Lyme and Bartonella treatment. More study is needed. I regret that we only examined cumanda for Bartonella and not Lyme.

Our conclusion was that cumunda hindered Bartonella more than Levofloxacin (levofloxacin), Zithromax (azithromycin), Rifabutin (mycobutin) and other proposed options. To determine treatment effect one needs to know **the indirect actions of Bartonella, Babesia, FL1953, Lyme, inflammation systems, etc. by lab analysis using different companies.**

Theophilus PA, Burugu D, Poururi A, Luecke DF, Sapi E. Effect of Medicinal Agents on the Different Forms of Borrelia burgdorferi Lyme disease or Lyme borreliosis is a tick-borne multisystemic disease caused by different species of Borrelia. <http://healthyats-nl.blogspot.com/2013/07/effect-of-medicinal-agents-stevia-and.html>

Erythromycin

Gomes found that erythromycin at low doses actually enhanced the growth of biofilms in *C. diphtheriae*. Penicillin acted the same way. Of further concern is that not only did these antibiotics increase biofilm formation but in this case they enhanced infections by strains of *C. diphtheriae*. Diphtheriae is a very dangerous infection without access to effective antibiotics. It is dangerous enough with good ones.

Returning to biofilm-promoted gum disease such as gingivitis, in the United States, over 50% of adults had gingivitis on an average of 3 to 4 teeth. Adult periodontitis, measured by the presence of periodontal pockets $> \text{ or } = 4 \text{ mm}$, was found in about 30% of the population on an average of 3 to 4 teeth. Lost gum attachment to teeth of at least 3 mm was found in 40% of the population (Oliver).

The density of adherent *P. gingivalis* cells were significantly decreased by using erythromycin at very low dosing called “sub-MIC levels.” One strain was not affected by erythromycin. Finally, erythromycin was not effective for inhibition of *P. gingivalis* biofilm cells at very low dosing.

Erythromycin Key Findings

- Low doses actually grew some biofilms
- Penicillin also grew some biofilms
- It enhanced strains of dangerous *C. diphtheriae*
- Gum disease from *P. gingivalis* cells was much less sticky at very low dosing.
- Erythromycin was not effective for inhibition of *P. gingivalis* biofilm cells at very low dosing.

Contacting Dr. Schaller

Should you wish to talk to Dr. Schaller he offers individualized education consults, which can be arranged by calling 239-263-0133. Please leave all your phone numbers, a working email and a fax number. These consults are typically in 15 minute units and can last as long as you wish. All that is required is the completion of a short informed consent form.

If you would like a full diagnostic consult or to see Dr. Schaller as a patient, know he treats patients from all over the USA and from outside the country. He meets with you first and then does follow-up care with you by phone.

If you would like to fly in to see Dr. Schaller, his staff are very familiar with all the closest airports, and we have special hotel discounts.